Harvard Pilgrim Health Care | TUFTS Health Plan

Pharmacy Medical Necessity Guidelines: **Simlandi® (adalimumab-ryvk)**

Effective: March 1, 2026

Guideline Type	☒ Prior Authorization ☐ Non-Formulary ☐ Step-Therapy ☐ Administrative

Applies to:

Commercial Products

☒ Harvard Pilgrim Health Care Commercial products; Fax: 617-673-0988

☒ Tufts Health Plan Commercial products; Fax: 617-673-0988

CareLink℠ – Refer to CareLink Procedures, Services and Items Requiring Prior Authorization

Public Plans Products

☒ Tufts Health Direct – A Massachusetts Qualified Health Plan (QHP) (a commercial product); Fax: 617-673-0988

Note: While you may not be the provider responsible for obtaining prior authorization, as a condition of payment you will need to ensure that prior authorization has been obtained.

Overview

Food and Drug Administration – Approved Indications

Simlandi (adalimumab-ryvk) is a tumor necrosis factor blocker indicated for:

Disease State	
Ankylosing Spondylitis	X
Crohn's Disease	X
Hidradenitis Suppurativa	X
Plaque Psoriasis	X
Polyarticular Juvenile Idiopathic Arthritis	X
Psoriatic Arthritis	X
Rheumatoid Arthritis	X
Ulcerative Colitis	X
Uveitis	X

Clinical Guideline Coverage Criteria

The plan may authorize coverage of Simlandi for Members when **ALL** of the following criteria are met:

Ankylosing Spondylitis

Initial Authorization Criteria

1. Documented diagnosis of active ankylosing spondylitis

AND

2. Patient is at least 18 years of age or older

AND

3. Prescribed by a rheumatologist

AND

4. Documentation of history of failure to **one (1)** prescription strength non-steroidal anti-inflammatory drug (e.g., celecoxib, diclofenac, ibuprofen, naproxen, meloxicam), unless contraindicated or clinically significant adverse effects are experienced *(Note: document drug, date and duration of trial)*

AND

5. Documentation the patient is not receiving the requested medication in combination with another biologic drug or with a targeted synthetic oral small molecule drug

Reauthorization Criteria

1. Documentation the patient has been established on the requested medication for at least 12 months

AND

2. Documented diagnosis of active ankylosing spondylitis

AND

3. Patient is at least 18 years of age or older

AND

4. Prescribed by a rheumatologist

AND

5. Documentation of clinically significant improvement or stabilization in clinical signs and symptoms of disease as determined by the prescriber

AND

6. Documentation the patient is not receiving the requested medication in combination with another biologic drug or with a targeted synthetic oral small molecule drug

Crohn's Disease

Initial Authorization Criteria

1. Documented diagnosis of moderately to severely active Crohn's disease

AND

2. Patient is at least 6 years of age or older

AND

3. Prescribed by a gastroenterologist

AND

4. Documentation of history of failure to **one (1)** of the following, unless contraindicated or clinically significant adverse effects are experienced *(Note: document drug, date, and duration of trial):*
    a. Corticosteroid
    b. 5-aminosalycylate
    c. 6-mercaptopurine
    d. Methotrexate

AND

5. Documentation the patient is not receiving the requested medication in combination with another biologic dug or with a targeted synthetic oral small molecule drug

Reauthorization Criteria

1. Documentation the patient has been established on the requested medication for at least 12 months

AND

2. Documented diagnosis of moderately to severely active Crohn's disease

AND

3. Patient is at least 6 years of age or older

AND

4. Prescribed by a gastroenterologist

AND

5. Documentation of clinically significant improvement or stabilization in clinical signs and symptoms of disease as determined by the prescriber

AND

6. Documentation the patient is not receiving the requested medication in combination with another biologic drug or with a targeted synthetic oral small molecule drug

Hidradenitis Suppurativa

Initial Authorization Criteria

1. Documented diagnosis of moderate to severe hidradenitis suppurativa

AND

2. Documentation of Hurley Stage II or III disease

AND

3. Patient is at least 12 years of age or older

AND

4. Prescribed by a dermatologist

AND

5. Documentation of history of failure to **three (3)** of the following conventional treatments, unless contraindicated or clinically significant adverse effects are experienced *(Note: document treatments, dates, and duration of trials)*

a. Local hygiene and ordinary hygiene
b. Weight reduction in patients who are obese
c. Use of ordinary soaps and antiseptic and antiperspirant agents (e.g., aluminum chloride hexahydrate) iv.
d. Application of warm compresses with sodium chloride solution or Burow's solution
e. Laser hair removal
f. Cessation of cigarette smoking
g. Medical anti-inflammatory or antiandrogen therapy such as oral or topical antibiotics, intralesional triamcinolone, spironolactone, or finasteride

AND

6. Documentation the patient is not receiving the requested medication in combination with another biologic drug or with a targeted synthetic oral small molecule drug

Reauthorization Criteria

1. Documentation the patient has been receiving the requested medication for at least six (6) months

AND

2. Documented diagnosis of moderate to severe hidradenitis suppurativa

AND

3. Patient is at least 12 years of age or older

AND

4. Prescribed by a dermatologist

AND

5. Submission of medical charts documenting the patient experienced a beneficial clinical response defined as an improvement from baseline in at least **one (1)** objective measure (e.g., Hurley staging, Sartorius score, Physician Global Assessment, Hidradenitis Suppurativa Severity Index)

AND

6. Submission of medical charts documenting that compared to baseline, the patient has experienced an improvement in at least one symptom of hidradenitis suppurativa such as decreased pain or drainage of lesions, nodules, or cysts

AND

7. Documentation the patient is not receiving the requested medication in combination with another biologic drug or with a targeted synthetic oral small molecule drug

Plaque Psoriasis

Initial Authorization Criteria

1. Documented diagnosis of moderate to severe chronic plaque psoriasis

AND

2. Documentation of **one (1)** of the following:
   a. Greater than or equal to 3% body surface area involvement
   b. Severe scalp psoriasis
   c. Palmoplantar, facial, or genital involvement

AND

3. Patient is at least 18 years of age or older

AND

4. Prescribed by a dermatologist

AND

5. Documentation of failure to a minimum 30-day supply (14-day supply for topical corticosteroids) to **one (1)** of the following topical therapies, unless contraindicated or clinically significant adverse effects are experienced: corticosteroid, vitamin D analog, tazarotene, calcineurin inhibitor *(Note: document drug, date and duration of trial)*

AND

6. Documentation the patient is not receiving the requested medication in combination with another biologic drug or with a targeted synthetic oral small molecule drug

Reauthorization Criteria

1. Documentation the patient has been receiving the requested medication for at least six (6) months

AND

2. Documented diagnosis of moderately to severe plaque psoriasis

AND

3. Patient is at least 18 years of age or older

AND

4. Prescribed by a dermatologist

AND

5. Submission of medical charts documenting the patient experienced a beneficial clinical response defined as reduction in body surface area affected by plaque psoriasis from baseline

AND

6. Submission of medical charts documenting that compared to baseline, the patient has experienced an improvement in at least one symptom of plaque psoriasis such as itching, redness, flaking, scaling, burning, cracking, or pain

AND

7. Documentation the patient is not receiving the requested medication in combination with another biologic drug or with a targeted synthetic oral small molecule drug

Polyarticular Juvenile Idiopathic Arthritis

Initial Authorization Criteria

1. Documented diagnosis of moderately to severely active polyarticular juvenile idiopathic arthritis

AND

2. Patient is at least 2 years of age or older

AND

3. Prescribed by a rheumatologist

AND

4. Documentation of history of failure to **one (1)** traditional disease modifying antirheumatic drug (e.g., methotrexate, leflunomide, hydroxychloroquine, sulfasalazine), unless contraindicated or clinically significant adverse effects are experienced *(Note: document drug, date, and duration of trial)*

AND

5. Documentation the patient is not receiving the requested medication in combination with another biologic drug or with a targeted synthetic oral small molecule drug

Reauthorization Criteria

1. Documentation the patient has been established on the requested medication for at least 12 months

AND

2. Documented diagnosis of active moderately to severely active polyarticular juvenile idiopathic arthritis

AND

3. Patient is at least 2 years of age or older

AND

4. Prescribed by a rheumatologist

AND

5. Documentation of clinically significant improvement or stabilization in clinical signs and symptoms of disease as determined by the prescriber

AND

6. Documentation the patient is not receiving the requested medication in combination with another biologic drug or with a targeted synthetic oral small molecule drug

Psoriatic Arthritis

Initial Authorization Criteria

1. Documented diagnosis of active psoriatic arthritis

AND

2. Documentation of **one (1)** of the following:
    a. Actively inflamed joints
    b. Dactylitis
    c. Enthesitis
    d. Axial disease
    e. Active skin and/or nail involvement

AND

3. Patient is at least 18 years of age or older

AND

4. Prescribed by a rheumatologist or dermatologist

AND

5. Documentation the patient is not receiving the requested medication in combination with another biologic drug or with a targeted synthetic oral small molecule drug

Reauthorization Criteria

1. Documentation the patient has been receiving the requested medication for at least six (6) months

AND

2. Documented diagnosis of active psoriatic arthritis

AND

3. Patient is at least 18 years of age or older

AND

4. Prescribed by a rheumatologist or dermatologist

AND

5. Submission of medical charts documenting the patient experienced a beneficial clinical response defined improvement from baseline in at least **one (1)** of the following:
    a. Body surface area affected by psoriatic arthritis
    b. Total active (swollen and tender) joint count

AND

6. Submission of medical charts documenting that compared to baseline, the patient has experienced an improvement in at least one symptom of psoriatic arthritis such as joint pain, morning stiffness, fatigue, joint or tendon sheath swelling

AND

7. Documentation the patient is not receiving the requested medication in combination with another biologic drug or with a targeted synthetic oral small molecule drug

Rheumatoid Arthritis

Initial Authorization Criteria

1. Documented diagnosis of moderately to severely active rheumatoid arthritis

AND

2. Patient is at least 18 years of age or older

AND

3. Prescribed by a rheumatologist

AND

4. Documentation of history of failure to **one (1)** traditional disease modifying antirheumatic drug (e.g., methotrexate, leflunomide, hydroxychloroquine, sulfasalazine), unless contraindicated or clinically significant adverse effects are experienced *(Note: document drug, date, and duration of trial)*

AND

5. Documentation the patient is not receiving the requested medication in combination with another biologic drug or with a targeted synthetic oral small molecule drug

Reauthorization Criteria

1. Documentation the Member has been established on the requested medication for at least 12 months

AND

2. Documented diagnosis of moderately to severely active rheumatoid arthritis

AND

3. Patient is at least 18 years of age or older

AND

4. Prescribed by a rheumatologist

AND

5. Documentation of clinically significant improvement or stabilization in clinical signs and symptoms of disease as determined by the prescriber

AND

6. Documentation the patient is not receiving the requested medication in combination with another biologic drug or with a targeted synthetic oral small molecule drug

Ulcerative Colitis

Initial Authorization Criteria

1. Documented diagnosis of moderately to severely active ulcerative colitis

AND

2. Patient is at least 5 years of age or older

AND

3. Prescribed by a gastroenterologist

AND

4. Documentation of history of failure to **one (1)** of the following, unless contraindicated or clinically significant adverse effects are experienced *(Note: document drug, date, and duration of trial):*
    a. Corticosteroid
    b. *5*-aminosalycyclate
    c. 6-mercaptopurine
    d. Methotrexate

AND

5. Documentation the patient is not receiving the requested medication in combination with another biologic drug or with a targeted synthetic oral small molecule drug

Reauthorization Criteria

1. Documentation the patient has been established on the requested medication for at least 12 months

AND

2. Documented diagnosis of moderately to severely active ulcerative colitis

AND

3. Patient is at least 5 years of age or older

AND

4. Prescribed by a gastroenterologist

AND

5. Documentation of clinically significant improvement or stabilization in clinical signs and symptoms of disease as determined by the prescriber

AND

6. Documentation the patient is not receiving the requested medication in combination with another biologic drug or with a targeted synthetic oral small molecule drug

Uveitis

Initial Authorization Criteria

1. Documented diagnosis of uveitis (non-infectious intermediate, posterior or panuveitis)

AND

2. Patient is at least 2 years of age or older

AND

3. Prescribed by an ophthalmologist, ocular immunologist, or rheumatologist

AND

4. Documentation of history of failure to conventional therapy (e.g., periocular, intraocular, or systemic corticosteroids; immunosuppressants), unless contraindicated or clinically significant adverse effects are experienced *(Note: document drug, date, and duration of trial)*

AND

5. Documentation the patient is not receiving the requested medication in combination with another biologic drug or with a targeted synthetic oral small molecule drug

Reauthorization Criteria

1. Documentation the patient has been established on the requested medication for at least 12 months

AND

2. Documented diagnosis of uveitis (non-infectious intermediate, posterior or panuveitis)

AND

3. Patient is at least 2 years of age or older

AND

4. Prescribed by an ophthalmologist, ocular immunologist, or rheumatologist

AND

5. Documentation of clinically significant improvement or stabilization in clinical signs and symptoms of disease as determined by the prescriber

AND

6. Documentation the patient is not receiving the requested medication in combination with another biologic drug or with a targeted synthetic oral small molecule drug

Limitations

1. Samples, free goods, or similar offerings of the requested medication do not qualify for an established clinical response and will not be considered for prior authorization.
2. Documentation of a Member being a social drinker does not qualify as a medically acceptable contraindication or clinical inappropriateness to methotrexate therapy.
3. For ankylosing spondylitis, Crohn's disease, polyarticular juvenile idiopathic arthritis, rheumatoid arthritis, ulcerative colitis and uveitis, authorizations will be provided in 12-month intervals. A patient who has received less than 12 months of therapy or who is restarting therapy with the requested drug is reviewed under Initial Authorization Criteria.
4. For hidradenitis suppurativa, plaque psoriasis and psoriatic arthritis, initial approval will be authorized for six (6) months. Reauthorization will be provided in 12-month intervals. A Member who has received less than six (6) months of therapy or who is restarting therapy with the requested drug is reviewed under Initial Authorization Criteria.
5. Members new the plan stable on the requested medication should be reviewed against Initial Authorization Criteria.

Codes

None

References

1. Simlandi (adalimumab-ryvk) [prescribing information]. Leesburg, VA: Alvotech USA Inc.;. October 2025.
2. Braun J, van den Berg R, Baraliakos X, et al. 2010 update of the ASAS/EULAR recommendations for the management of Ankylosing spondylitis. Ann Rheum Dis. 2011;70:896-904.
3. Felson D, Anderson J, Boers M, et al. American College of Rheumatology preliminary definition of improvement in rheumatoid arthritis. Arthritis Rheum 1995; 38:727-35.
4. Mease P, Gladman D, Ritchlin C, et al. Adalimumab for the treatment of patients with moderately to severely active psoriatic arthritis. Arthritis and Rheumatism 2005; 52:3279-89.
5. Braun J, Davis J et al. First update of the international ASAS consensus statement for the use of anti-TNF agents in patients with ankylosing spondylitis. Ann Rheum Dis. 2006; 65(3):316-20.
6. Hanauer SB, Sandborn WJ, Rutgeerts P, et al. Human Anti-Tumor Necrosis Factor Monoclonal Antibody (Adalimumab) in Crohn's Disease: the CLASSIC-I Trial. Gastroenterology. 2006; 130: 323-333.
7. Colombel JF, Sandborn WJ, Rutgeerts P, et al. Adalimumab for Maintenance of Clinical Response and Remission in Patients With Crohn's Disease: The CHARM Trial. Gastroenterology. 2007; 132: 53-65.
8. Menter A, Strober BE, Kaplan DH, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics. J Am Acad Dermatol 2019;80:1029-72.
9. Singh JA, Guyatt G, Ogdie A, et al. 2018 American College of Rheumatology/National Psoriasis Foundation guideline for the treatment of psoriatic arthritis. Arthritis Rheumatol. 2019;71(1):5-32.
10. Lichtenstein GR, Loftus EV, Isaacs KL, et al. ACG clinical guideline: management of Crohn's disease in adults. Am J Gastroenterol. 2018;113:481-517.
11. Singh S, Loftus EV, Jr, Limketkai BN, et al. AGA Living Clinical Practice Guideline on Pharmacological Management of Moderate-to-Severe Ulcerative Colitis. Gastroenterology. 2024 Dec;167(7):1307-1343.
12. Fraenkel L, Bathon JM, England BR, et al. 2021 American College of Rheumatology Guideline for the Treatment of Rheumatoid Arthritis. Arthritis Care Res. 2021;73(7):924-939.
13. Ringold S, Angeles-Han ST, Beukelman T, et al. 2019 American College of Rheumatology/Arthritis Foundation guideline for the treatment of juvenile idiopathic arthritis: therapeutic approaches for non-systemic polyarthritis, sacroiliitis, and enthesitis. Arthritis Rheumatol. 2019;71(6):846-863.
14. Rosenbaum JT. Uveitis: Treatment. In: Romain PL, ed. UpToDate [database online]. Waltham, MA: UpToDate, 2021. https://www.uptodate.com. Accessed October 20, 2021.
15. Ingram JR. Hidradenitis suppurativa: Management. In: Ofori AO, ed. UpToDate [database online]. Waltham, MA: UpToDate, 2021. https://www.uptodate.com. Accessed October 20, 2021.

Approval And Revision History

September 13, 2022: Reviewed by the Pharmacy & Therapeutics Committee.

Subsequent endorsement date(s) and changes made:

- December 12, 2023: No changes
- November 12, 2024: No changes (eff 11/12/24)
- October 14, 2024: Provider requirements no longer allow with in consultation with the specialist. Added coverage criteria to prevent duplicative targeted immunomodulator therapy. For hidradenitis suppurativa, plaque psoriasis and psoriatic arthritis, initial approvals limited to 6 months with annual reauthorization coverage criteria added, added disease severity requirements, updated Initial Authorization Criteria prerequisite language (no change in prerequisites) and removed previous treatment with a biologic agent indicated for the requested use and Humira stability language. For ankylosing spondylitis,

Crohn's disease, polyarticular juvenile idiopathic arthritis, rheumatoid arthritis, ulcerative colitis and uveitis, approvals limited to 12 months with reauthorization coverage criteria added, updated Initial Authorization Criteria prerequisite language (no change in prerequisites), and removed previous treatment with a biologic agent indicated for the requested use and Humira stability language. (eff 1/1/26).

- December 9, 2025: Simlandi moved from non-preferred to preferred status. Humira moved from preferred to non-preferred status. Medical Necessity Guideline updated from Humira to Simlandi (eff 3/1/26).

Background, Product and Disclaimer Information

Pharmacy Medical Necessity Guidelines have been developed for determining coverage for plan benefits and are published to provide a better understanding of the basis upon which coverage decisions are made. The plan makes coverage decisions on a case-by-case basis considering the individual member's health care needs. Pharmacy Medical Necessity Guidelines are developed for selected therapeutic classes or drugs found to be safe, but proven to be effective in a limited, defined population of patients or clinical circumstances. They include concise clinical coverage criteria based on current literature review, consultation with practicing physicians in the service area who are medical experts in the particular field, FDA and other government agency policies, and standards adopted by national accreditation organizations. The plan revises and updates Pharmacy Medical Necessity Guidelines annually, or more frequently if new evidence becomes available that suggests needed revisions.

For self-insured plans, coverage may vary depending on the terms of the benefit document. If a discrepancy exists between a Pharmacy Medical Necessity Guideline and a self-insured Member’s benefit document, the provisions of the benefit document will govern.

Treating providers are solely responsible for the medical advice and treatment of members. The use of this policy is not a guarantee of payment or a final prediction of how specific claim(s) will be adjudicated. Claims payment is subject to member eligibility and benefits on the date of service, coordination of benefits, referral/authorization and utilization management guidelines when applicable, and adherence to plan policies and procedures and claims editing logic.