Pharmacy Medical Necessity Guidelines:
Wainua (eplontersen)

Effective: July 14, 2026

Prior Authorization Required	√	Type of Review – Care Management	
Not Covered		Type of Review – Clinical Review	√
Pharmacy (RX) or Medical (MED) Benefit	RX	Department to Review	RXUM
These pharmacy medical necessity guidelines apply to the following: ☒ Tufts Health RITogether – A Rhode Island Medicaid Plan		**Fax Numbers:** RXUM: 617.673.0939	

Note: This guideline does not apply to Medicare Members (includes dual eligible Members).

OVERVIEW

FOOD AND DRUG ADMINISTRATION-APPROVED INDICATIONS

Wainua (eplontersen) is a transthyretin-directed antisense oligonucleotide indicated for treatment of the polyneuropathy of hereditary transthyretin-mediated amyloidosis in adults.

COVERAGE GUIDELINES

The plan may authorize coverage of **Wainua** (eplontersen) for Members, when the following criteria are met:

Initial Authorization Criteria

1. Documented diagnosis of hereditary transthyretin-mediated amyloidosis with polyneuropathy

AND

2. Documentation in the Member's medical record of transthyretin (TTR) genetic mutation

AND

3. The patient is at least 18 years of age or older

AND

4. Prescribed by or in consultation with a neurologist or a provider who specializes in amyloidosis

AND

5. Documentation the patient is currently experiencing signs and symptoms of polyneuropathy defined as **one (1)** of the following:
    a. Baseline polyneuropathy disability (PND) score less than or equal to IIIb
    b. Baseline familial amyloid polyneuropathy Stage 1 or 2
    c. Baseline neuropathy impairment score between 10 and 130

AND

6. Documentation the patient has not had a prior liver transplant

AND

7. Documentation the requested medication will not be used concomitantly with another medication indicated for the management of cardiomyopathy or neuropathy of transthyretin-mediated amyloidosis (e.g., Amvuttra, Onpattro, Vyndamax)

Reauthorization Criteria

1. Documented diagnosis of hereditary transthyretin-mediated amyloidosis with polyneuropathy

AND

2. Documentation in the Member's medical record of transthyretin (TTR) genetic mutation

AND

3. The patient is at least 18 years of age or older

AND

4. Prescribed by or in consultation with a neurologist or a provider who specializes in amyloidosis

AND

5. Documentation the requested medication will not be used concomitantly with another medication indicated for the management of cardiomyopathy or neuropathy of transthyretin-mediated amyloidosis (e.g., Amvuttra, Onpattro, Vyndamax)

AND

6. Documentation of a positive clinical response to therapy as evidenced by improved or stable disease as defined by **one (1)** of the following scales:

a. Modified Neuropathy Impairment Scale+7 composite scale
b. Norfolk Quality of Life-Diabetic Neuropathy total score

LIMITATIONS

- Initial coverage of Wainua will be authorized for 12 months. Reauthorization of Wainua will be provided in 12-month intervals.
- Members new to the plan stable on Wainua should be reviewed against Reauthorization criteria.
- For a non-formulary medication request, please refer to the Pharmacy Medical Necessity Guidelines for Formulary Exceptions and submit a formulary exception request to the plan as indicated.

CODES

None

REFERENCES

1. Adams D, Gonzalez-Duarte A, O'Riordan WD, et al. Patisiran, an RNAi therapeutic, for hereditary transthyretin amyloidosis. N Engl J Med. 2018;379(1):11-21.
2. Ando Y, Coelho T, Berk JL, et al. Guideline of transthyretin-related hereditary amyloidosis for clinicians. Orphanet Journal of Rare Diseases. 2013;8(31):1-18.
3. Benson MD, Waddington-Cruz M, Berk J, et al. Inotersen treatment for patients with hereditary transthyretin amyloidosis. N Engl J Med. 2018;379(1):22-31.
4. Brannagan T, Wang AK, Coelho T, et al. Open label extension of the phase 3 study NEURO-TTR to assess the long-term efficacy and safety of inotersen in patients with hereditary transthyretin amyloidosis. Neurology. 2018;90(15 Suppl). Abstract P1.324.
5. Lasser KE, Mickle K, Chapman R, et al. Inotersen and patisiran for hereditary transthyretin amyloidosis: effectiveness and value. Evidence report. 2018 August 29. Available from Internet. Accessed 2018 September 12.
6. Wainua (eplontersen) [package insert]. Wilmington, DE: AstraZeneca Pharmaceuticals LP.; Sep 2024.

APPROVAL HISTORY

June 11, 2024: Reviewed by Pharmacy & Therapeutics Committee.
July 8, 2025: No changes (eff 7/8/25)
July 14, 2026: No changes (eff 7/14/26)

BACKGROUND, PRODUCT AND DISCLAIMER INFORMATION

Pharmacy Medical Necessity Guidelines have been developed for determining coverage for plan benefits and are published to provide a better understanding of the basis upon which coverage decisions are made. The plan makes coverage decisions on a case-by-case basis considering the individual member's health care needs. Pharmacy Medical Necessity Guidelines are developed for selected therapeutic classes or drugs found to be safe, but proven to be effective in a limited, defined population of patients or clinical circumstances. They include concise clinical coverage criteria based on current literature review, consultation with practicing physicians in the service area who are medical experts in the particular field, FDA and other government agency policies, and standards adopted by national accreditation organizations. The plan revises and updates Pharmacy Medical Necessity Guidelines annually, or more frequently if new evidence becomes available that suggests needed revisions.

For self-insured plans, coverage may vary depending on the terms of the benefit document. If a discrepancy exists between a Pharmacy Medical Necessity Guideline and a self-insured Member's benefit document, the provisions of the benefit document will govern.

Treating providers are solely responsible for the medical advice and treatment of members. The use of this policy is not a guarantee of payment or a final prediction of how specific claim(s) will be adjudicated. Claims payment is subject to member eligibility and benefits on the date of service, coordination of benefits, referral/authorization and utilization management guidelines when applicable, and adherence to plan policies and procedures and claims editing logic.

Provider Services